

objects, that is that ' $\Box F$ ', if true of an object, is true of it regardless of the way in which it is referred to, then one should better not quantify into causal contexts; one should avoid contrary-to-fact conditionals, scientific law-statements, confirmation statements, and many types of probability statements and disposition terms—if one wants to make sense.

V

SEMANTICAL CONSIDERATIONS ON
MODAL LOGIC

SAUL A. KRIPKE

This paper gives an exposition of some features of a semantical theory of modal logics.¹ For a certain quantified extension of *SS*, this theory was presented in 'A Completeness Theorem in Modal Logic',² and it has been summarized in 'Semantical Analysis of Modal Logic'.³ The present paper will concentrate on one aspect of the theory—the introduction of quantifiers—and it will restrict itself in the main to one method of achieving this end. The emphasis of the paper will be purely semantical, and hence it will omit the use of semantic tableaux, which is essential to a full presentation of the theory.⁴ Proofs, also, will largely be suppressed.

We consider four modal systems. Formulae A, B, C, \dots are built out of atomic formulae P, Q, R, \dots , using the connectives $\wedge, \sim,$ and \Box . The system M has the following axiom schemes and rules:

A0. Truth-functional tautologies

A1. $\Box A \supset A$

A2. $\Box(A \supset B) \supset \Box A \supset \Box B$

R1. $A, A \supset B/B$

R2. $A/\Box A$

If we add the following axiom scheme, we get *S4*:

$\Box A \supset \Box \Box A$

From *Acta Philosophica Fennica*, 16 (1963), 83–94. Reprinted by permission of the author and the publishers, Societas Philosophica Fennica, Helsinki.

¹ The theory given here has points of contact with many authors: For lists of these, see S. Kripke, 'Semantical Analysis of Modal Logic', *Zeitschrift für Mathematische Logik und Grundlagen der Mathematik*, 9 (1963), 67–96, and J. Hintikka, 'Modality and Quantification' *Theoria*, 27 (1961) 119–28. The authors closest to the present theory appear to be Hintikka and Kanger. The present treatment of quantification, however, is unique as far as I know, although it derives some inspiration from acquaintance with the very different methods of Prior and Hintikka.

² *Journal of Symbolic Logic*, 24 (1959), 1–15.

³ *Ibid.*, pp. 323–4 (Abstract).

⁴ For these see 'A Completeness Theorem in Modal Logic', *Journal of Symbolic Logic*, 24 (1959), 1–15 and 'Semantical Analysis of Modal Logic', *Zeitschrift für Mathematische Logik und Grundlagen der Mathematik*, 9, 67–96.

We get the *Brouwersche* system if we add to *M*:

$$\begin{aligned} A &\supset \Box \Diamond A \\ \text{S5, if we add:} \\ \Diamond A &\supset \Box \Diamond A \end{aligned}$$

Modal systems whose theorems are closed under the rules R1 and R2, and include all theorems of *M*, are called 'normal'. Although we have developed a theory which applies to such non-normal systems as Lewis's S2 and S3, we will restrict ourselves here to normal systems.

To get a semantics for modal logic, we introduce the notion of a (normal) *model structure*. A model structure (m.s.) is an ordered triple (G, K, R) where K is a set, R is a reflexive relation on K , and $G \in K$. Intuitively, we look at matters thus: K is the set of all 'possible worlds'; G is the 'real world'. If H_1 and H_2 are two worlds, $H_1 R H_2$ means intuitively that H_2 is 'possible relative to' H_1 ; i.e., that every proposition true in H_2 is possible in H_1 . Clearly, then, the relation R should indeed be reflexive; every world H is possible relative to itself, since every proposition true in H is, *a fortiori*, possible in H . Reflexivity is thus an intuitively natural requirement. We may impose additional requirements, corresponding to various 'reduction axioms' of modal logic: If R is transitive, we call (G, K, R) an S4-m.s.; if R is symmetric, (G, K, R) is a *Brouwersche* m.s.; and if R is an equivalence relation, we call (G, K, R) an S5-m.s. A model structure without restriction is also called an *M*-model structure.

To complete the picture, we need the notion of *model*. Given a model structure (G, K, R) , a *model* assigns to each atomic formula (propositional variable) P a truth-value T or F in each world $H \in K$. Formally, a *model* φ on a m.s. (G, K, R) is a binary function $\varphi(P, H)$, where P varies over atomic formulae and H varies over elements of K , whose range is the set $\{T, F\}$. Given a model, we can define the assignments of truth-values to non-atomic formulae by induction. Assume $\varphi(A, H)$ and $\varphi(B, H)$ have already been defined for all $H \in K$. Then if $\varphi(A, H) = \varphi(B, H) = T$, define $\varphi(A \wedge B, H) = T$; otherwise, $\varphi(A \wedge B, H) = F$. $\varphi(\sim A, H)$ is defined to be F iff $\varphi(A, H) = T$; otherwise, $\varphi(\sim A, H) = T$. Finally, we define $\varphi(\Box A, H) = T$ iff $\varphi(A, H') = T$ for every $H' \in K$ such that $H R H'$; otherwise, $\varphi(\Box A, H) = F$. Intuitively, this says that A is necessary in H iff A is true in all worlds H' possible relative to H .

Completeness theorem. $\vdash A$ in *M* (S4, S5, the *Brouwersche* system) if and only if $\varphi(A, G) = T$ for every model φ on an *M*-(S4-, S5-, *Brouwersche*) model structure (G, K, R) .⁵

⁵ For a proof, see 'Semantical Analysis . . .', *Zeitschrift* . . ., 9.

This completeness theorem equates the syntactical notion of *provability* in a modal system with a semantical notion of *validity*.

The rest of this paper concerns, with the exception of some concluding remarks, the introduction of quantifiers. To do this, we must associate with each world a domain of individuals, the individuals that exist in that world. Formally, we define a *quantificational model structure* (q.m.s.) as a model structure (G, K, R) , together with a function ψ which assigns to each $H \in K$ a set $\psi(H)$, called the *domain* of H . Intuitively $\psi(H)$ is the set of all individuals existing in H . Notice, of course, that $\psi(H)$ need not be the same set for different arguments H , just as, intuitively, in worlds other than the real one, some actually existing individuals may be absent, while new individuals, like Pegasus, may appear.

We may then add, to the symbols of modal logic, an infinite list of individual variables x, y, z, \dots , and, for each nonnegative integer n , a list of n -adic predicate letters P^n, Q^n, \dots , where the superscripts will sometimes be understood from the context. We count propositional variables (atomic formulae) as '0-adic' predicate letters. We then build up well-formed formulae in the usual manner, and can now prepare ourselves to define a *quantificational model*.

To define a *quantificational model*, we must extend the original notion, which assigned a truth-value to each atomic formula in each world. Analogously, we must suppose that in each world a given n -adic predicate letter determines a certain set of ordered n -tuples, its *extension* in that world. Consider, for example, the case of a monadic predicate letter $P(x)$. We would like to say that, in the world H , the predicate $P(x)$ is true of some individuals in $\psi(H)$ and false of others; formally, we would say that, relative to certain assignments of elements of $\psi(H)$ to x , $\varphi(P(x), H) = T$ and relative to others $\varphi(P(x), H) = F$. The set of all individuals of which P is true is called the *extension* of P in H . But there is a problem: should $\varphi(P(x), H)$ be given a truth-value when x is assigned a value in the domain of some other world H' and not in the domain of H ? Intuitively, suppose $P(x)$ means 'x is bald'—are we to assign a truth-value to the substitution instance 'Sherlock Holmes is bald'? Holmes does not exist, but in other states of affairs, he would have existed. Should we assign a definite truth-value to the statement that he is bald, or not? Frege⁶ and Strawson⁷

⁶ G. Frege 'Über Sinn und Bedeutung', *Zeitschrift für Philosophie und philosophische Kritik*, 100 (1892), 25–50. English translations in Geach and Black, *Translations from the Philosophical Writings of Gottlob Frege*, (Oxford: Blackwell, 1952), and in Feigl and Sellars (eds.), *Readings in Philosophical Analysis* (New York: Appleton Century Crofts, 1949).

⁷ P. F. Strawson, 'On referring', *Mind*, n.s., 59 (1950), 320–44.

would not assign the statement a truth-value; Russell would.⁸ For the purposes of modal logic we hold that different answers to this question represent alternative *conventions*. All are tenable. The only existing discussions of this problem I have seen—those of Hintikka⁹ and Prior¹⁰—adopt the Frege-Strawson view. This view necessarily must lead to some modification of the usual modal logic. The reason is that the semantics for modal propositional logic, which we have already given, assumed that every formula must take a truth-value in each world; and now, for a formula $A(x)$ containing a free variable x , the Frege-Strawson view requires that it not be given a truth-value in a world H when the variable x is assigned an individual not in the domain of that world. We thus can no longer expect that the original laws of modal propositional logic hold for statements containing free variables, and are faced with an option: either revise modal propositional logic or restrict the rule of substitution. Prior does the former, Hintikka the latter. There are further alternatives the Frege-Strawson choice involves: Should we take $\Box A$ (in H) to mean that A is *true* in all possible worlds (relative to H), or just *not false* in any such world? The second alternative merely demands that A be either true or lack a truth-value in each world. Prior, in his system Q , in effect admits both types of necessity, one as ' L ' and the other as ' NMN '. A similar question arises for conjunction: if A is false and B has no truth-value, should we take $A \wedge B$ to be false or truth-valueless?

In a full statement of the semantical theory, we would explore all these variants of the Frege-Strawson view. Here we will take the other option, and assume that a statement containing free variables has a truth-value in each world for every assignment to its free variables.¹¹ Formally, we state the matter as follows: Let $U = \prod_{H \in K} \psi(H)$. U^n is the n th Cartesian product of U with itself. We define a quantificational model on a q.m.s. (G, K, R) as a binary function $\varphi(P^n, H)$, where the first variable ranges over n -adic predicate letters, for arbitrary n , and H ranges

⁸ Bertrand Russell, 'On denoting', *Mind*, n.s., 14 (1905), 479–93.

⁹ 'Modality and Quantification'.

¹⁰ A. N. Prior, *Time and Modality* (Oxford: Clarendon Press, 1957, viii+148 pp.).

¹¹ It is natural to assume that an atomic predicate should be false in a world H of all those individuals not existing in that world; that is, that the extension of a predicate letter must consist of actually existing individuals. We can do this by requiring semantically that $\varphi(P^n, H)$ be a subset of $[\psi(H)]^n$; the semantical treatment below would otherwise suffice without change. We would have to add to the axiom system below all closures of formulae of the form $P^n(x_1, \dots, x_n) \wedge (y)A(y) \supset A(x_i)$ ($1 \leq i \leq n$). We have chosen not to do this because the rule of substitution would no longer hold; theorems would hold for atomic formulae which would not hold when the atomic formulae are replaced by arbitrary formulae. (This answers a question of Putnam and Kalmar.)

over elements of K . If $n = 0$, $\varphi(P^n, H) = T$ or F ; if $n \geq 1$, $\varphi(P^n, H)$ is a subset of U^n . We now define, inductively, for every formula A and $H \in K$, a truth-value $\varphi(A, H)$, relative to a given assignment of elements of U to the free variables of A . The case of a propositional variable is obvious. For an atomic formula $P^n(x_1, \dots, x_n)$, where P^n is an n -adic predicate letter and $n \geq 1$, given an assignment of elements a_1, \dots, a_n of U to x_1, \dots, x_n , we define $\varphi(P^n(x_1, \dots, x_n), H) = T$ if the n -tuple (a_1, \dots, a_n) is a member of $\varphi(P^n, H)$; otherwise, $\varphi(P^n(x_1, \dots, x_n), H) = F$, relative to the given assignment. Given these assignments for atomic formulae, we can build up the assignments for complex formulae by induction. The induction steps for the propositional connectives \wedge, \sim, \Box , have already been given. Assume we have a formula $A(x, y_1, \dots, y_n)$, where x and the y_i are the only free variables present, and that a truth-value $\varphi(A(x, y_1, \dots, y_n), H)$ has been defined for each assignment to the free variables of $A(x, y_1, \dots, y_n)$. Then we define $\varphi((x)A(x, y_1, \dots, y_n), H) = T$ relative to an assignment of b_1, \dots, b_n to y_1, \dots, y_n (where the b_i are elements of U), if $\varphi(A(x, y_1, \dots, y_n), H) = T$ for every assignment of a, b_1, \dots, b_n to x, y_1, \dots, y_n , respectively, where $a \in \psi(H)$; otherwise, $\varphi((x)A(x, y_1, \dots, y_n), H) = F$ relative to the given assignment. Notice that the restriction $a \in \psi(H)$ means that, in H , we quantify only over the objects actually existing in H .

To illustrate the semantics, we give counterexamples to two familiar proposals for laws of modal quantification theory—the 'Barcan formula' $(x)\Box A(x) \supset \Box(x)A(x)$ and its converse $\Box(x)A(x) \supset (x)\Box A(x)$. For each we consider a model structure (G, K, R) , where $K = \{G, H\}$, $G \neq H$, and R is simply the Cartesian product K^2 . Clearly R is reflexive, transitive, and symmetric, so our considerations apply even to S5.

For the Barcan formula, we extend (G, K, R) to a quantificational model structure by defining $\psi(G) = \{a\}$, $\psi(H) = \{a, b\}$, where a and b are distinct. We then define, for a monadic predicate letter P , a model φ in which $\varphi(P, G) = \{a\}$, $\varphi(P, H) = \{a\}$. Then clearly $\Box P(x)$ is true in G when x is assigned a ; and since a is the only object in the domain of G , so is $(x)\Box P(x)$. But, $(x)P(x)$ is clearly false in H (for $\varphi(P(x), H) = F$ when x is assigned b), and hence $\Box(x)P(x)$ is false in G . So we have a counterexample to the Barcan formula. Notice that this counterexample is quite independent of whether $P(x)$ is assigned a truth-value in G or not when x is assigned δ , so also it applies to the systems of Hintikka and Prior. Such counterexamples can be disallowed, and the Barcan formula reinstated, only if we require a model structure to satisfy the condition that $\psi(H') \subseteq \psi(H)$ whenever HRH' ($H, H' \in K$).

For the converse of the Barcan formula, set $\psi(G) = \{a, b\}$, $\psi(H) = \{a\}$,

where again $a \neq b$. Define $\varphi(P, G) = \{a, b\}$, $\varphi(P, H) = \{a\}$, where P is a given monadic predicate letter. Then clearly $(x)P(x)$ holds in both G and H , so that $\varphi(\Box(x)P(x), G) = T$. But $\varphi(P(x), H) = F$ when x is assigned b , so that, when x is assigned b , $\varphi(\Box P(x), G) = F$. Hence $\varphi(\Box(x)\Box P(x), G) = F$, and we have the desired counterexample to the converse of the Barcan formula. This counterexample, however, depends on asserting that, in H , $P(x)$ is actually *false* when x is assigned b ; it might thus disappear if, for this assignment, $P(x)$ were declared to lack truth-value in H . In this case, we will still have a counterexample if we require a necessary statement to be *true* in all possible worlds (Prior's 'L'), but not if we merely require that it never be false (Prior's 'NMN'). On our present convention, we can eliminate the counterexample only by requiring, for each q.m.s., that $\psi(H) \subseteq \psi(H')$ whenever HRH' .

These counterexamples lead to a peculiar difficulty: We have given countermodels, in quantified S5, to both the Barcan formula and its converse. Yet Prior appears to have shown¹² that the Barcan formula is derivable in quantified S5; and the converse seems derivable even in quantified M by the following argument:

- (A) $(x)A(x) \supset A(y)$ (by quantification theory)
- (B) $\Box((x)A(x) \supset A(y))$ (by necessitation)
- (C) $\Box((x)A(x) \supset A(y)) \supset \Box(x)A(x) \supset \Box A(y)$ (Axiom A2)
- (D) $\Box(x)A(x) \supset \Box A(y)$ (from (B) and (C))
- (E) $(y)(\Box(x)A(x) \supset \Box A(y))$ (generalizing on (D))
- (F) $\Box(x)A(x) \supset (y)\Box A(y)$ (by quantification theory, and (E))

We seem to have derived the conclusion using principles that should all be valid in the model-theory. Actually, the flaw lies in the application of necessitation to (A). In a formula like (A), we give the free variables the

¹² See 'Modality and Quantification in s5', *Journal of Symbolic Logic*, 21 (1956), 60-2.

¹³ It is not asserted that the generality interpretation of theorems with free variables is the only possible one. One might wish a formula A to be provable iff, for each model φ , $\varphi(A, G) = T$ for every assignment to the free variables of A . But then $(x)A(x) \supset A(y)$ will not be a theorem; in fact, in the countermodel above to the Barcan formula, $\varphi((x)P(x) \supset P(y), G) = F$ if y is assigned b . Thus quantification theory would have to be revised along the lines proposed by Hintikka (in 'Existential Presuppositions and Existential Commitments', *Journal of Philosophy*, 56 (1959), 125-37) and by H. Leblanc and T. Hailperin (in 'Nondesignating Singular Terms', *Philosophical Review*, 68 (1959), 239-43). This procedure has much to recommend it, but we have not adopted it since we wished to show that the difficulty can be solved without revising quantification theory or modal propositional logic.

generality interpretation:¹³ When (A) is asserted as a theorem, it abbreviates assertion of its ordinary universal closure

$$(A') (y)((x)A(x) \supset A(y))$$

Now if we applied necessitation to (A'), we would get

$$(B') \Box(y)((x)A(x) \supset A(y))$$

On the other hand, (B) itself is interpreted as asserting

$$(B'') (y)\Box((x)A(x) \supset A(y))$$

To infer (B'') from (B'), we would need a law of the form $\Box(y)C(y) \supset (y)\Box C(y)$, which is just the converse Barcan formula that we are trying to prove. In fact, it is readily checked that (B'') fails in the countermodel given above for the converse Barcan formula, if we replace $A(x)$ by $P(x)$.

We can avoid this sort of difficulty if, following Quine¹⁴ we formulate quantification theory so that only *closed* formulae are asserted. Assertion of formulae containing free variables is at best a convenience; assertion of $A(x)$ with free x can always be replaced by assertion of $(x)A(x)$.

If A is a formula containing free variables, we define a *closure* of A to be any formula without free variables obtained by prefixing universal quantifiers and necessity signs, in any order, to A . We then define the axioms of quantified M to be the closures of the following schemata:

- (0) Truth-functional tautologies
- (1) $\Box A \supset A$
- (2) $\Box(A \supset B) \supset \Box A \supset \Box B$
- (3) $A \supset (x)A$, where x is not free in A
- (4) $(x)(A \supset B) \supset (x)A \supset (x)B$
- (5) $(y)((x)A(x) \supset A(y))$

The rule of inference is detachment for material implication. Necessitation can be obtained as a derived rule.

To obtain quantified extensions of S4, S5, the *Brouwersche* system, simply add to the axiom schemata all closures of the appropriate reduction axiom.

The systems we have obtained have the following properties: They are a straightforward extension of the modal propositional logics, without the modifications of Prior's Q; the rule of substitution holds without restriction, unlike Hintikka's presentation; and nevertheless neither the Barcan formula nor its converse is derivable. Further, all the laws of quantification theory—modified to admit the empty domain—hold.

¹⁴ W. Quine, *Mathematical Logic* (Cambridge, Mass.: Harvard Univ. Press, 1940; 2nd edn., rev., 1951, xii+346 pp.).

The semantical completeness theorem we gave for modal propositional logic can be extended to the new systems.

We can introduce *existence as a predicate* in the present system if we like. Semantically, existence is a monadic predicate $E(x)$ satisfying, for each model φ on a m.s. (G, K, R) , the identity $\varphi(E, H) = \psi(H)$ for every $H \in K$. Axiomatically, we can introduce it through the postulation of closures of formulae of the form: $(x)A(x) \wedge E(y) \supset A(y)$, and $(x)E(x)$. The predicate P used above in the counter-example to the converse Barcan formula can now be recognized as simply existence. This fact shows how existence differs from the tautological predicate $A(x) \vee \sim A(x)$ even though $\Box(x)E(x)$ is provable. For although $(x)\Box(A(x) \vee \sim A(x))$ is valid, $(x)\Box E(x)$ is not; although it is necessary that every thing exists, it does not follow that everything has the property of necessary existence.

We can introduce identity semantically in the model theory by defining $x = y$ to be true in a world H when x and y are assigned the same value and otherwise false; existence could then be defined in terms of identity, by stipulating that $E(x)$ means $(\exists y)(x = y)$. For reasons not given here, a broader theory of identity could be obtained if we complicated the notion of quantificational model structure.

We conclude with some brief and sketchy remarks on the 'provability' interpretations of modal logics, which we give in each case for propositional calculus only. The reader will have obtained the main point of this paper if he omits this section. Provability interpretations are based on a desire to adjoin a necessity operator to a formal system, say Peano arithmetic, in such a way that, for any formula A of the system, $\Box A$ will be interpreted as true iff A is provable in the system. It has been argued that such 'provability' interpretations of a model operator are dispensable in favour of a provability *predicate*, attaching to the Gödel number of A ; but Professor Montague's contribution to the present volume casts at least some doubt on this viewpoint.

Let us consider the formal system PA of Peano arithmetic, as formalized in Kleene.¹⁵ We adjoin to the formation rules operators \wedge , \sim , and \Box (the conjunction and negation adjoined are to be distinct from those of the original system), operating on closed formulae only. In the model theory we gave above, we took atomic formulae to be propositional variables, or predicate letters followed by parenthesized individual variables; here we take them to be simply the closed well-formed formulae of PA (not just the atomic formulae of PA). We define a model structure (G, K, R) , where K is the set of all distinct (non-isomorphic) countable models of

¹⁵ S. C. Kleene, *Introduction to Metamathematics* (New York: D. Van Nostrand, 1952, x+550 pp.).

PA, G is the standard model in the natural numbers, and R is the Cartesian product K^2 . We define a model φ by requiring that, for any atomic formula P and $H \in K$, $\varphi(P, H) = T(F)$ iff P is true (false) in the model H . (Remember, P is a wff of PA, and H is a countable model of PA.) We then build up the evaluation for compound formulae as before.¹⁶ To say that A is true is to say it is true in the real world G ; and, for any atomic P , $\varphi(\Box P, G) = T$ iff P is provable in PA. (Notice that $\varphi(P, G) = T$ iff P is true in the intuitive sense). Since (G, K, R) is an S5-m.s., all the laws of S5 will be valid on this interpretation; and we can show that *only* the laws of S5 are generally valid. (For example, if P is Gödel's undecidable formula, $\varphi(\Box P \vee \Box \sim P, G) = F$, which is a counterexample to the 'law' $\Box A \vee \Box \sim A$.)

Another provability interpretation is the following: Again we take the atomic formulae to be the closed wffs of PA, and then build up new formulae using the adjoined connectives \wedge , \sim , and \Box . Let K be the set of all ordered pairs (E, α) , where E is a consistent extension of PA, and α is a (countable) model of the system E . Let $G = (PA, \alpha_0)$, where α_0 is the standard model of PA. We say $(E, \alpha) R (E', \alpha')$, where (E, α) and (E', α') are in K , iff E' is an extension of E . For atomic P , define $\varphi(P, (E, \alpha)) = T(F)$ iff P is true (false) in α . Then we can show, for atomic P , that $\varphi(\Box P, (E, \alpha)) = T$ iff P is provable in E ; in particular, $\varphi(\Box P, G) = T$ iff P is provable in PA. Since (G, K, R) is an S4-m.s., all the laws of S4 hold. But not all the laws of S5 hold; if P is Gödel's undecidable formula, $\varphi((\sim \Box P \supset \Box \sim \Box P), G) = F$. But some laws are valid which are not provable in S4; in particular, we can prove for any A , $\varphi(\Box \sim \Box(\Diamond A \wedge \Diamond \sim A), G) = T$, which yields the theorems of McKinsey's S4.1.¹⁷ By suitable modifications this difficulty could be removed; but we do not go into the matter here.

Similar interpretations of M and the *Brouwersche* system could be stated; but, in the present writer's opinion, they have less interest than those given above. We mention one more class of provability interpretations, the 'reflexive' extensions of PA. Let E be a formal system containing

¹⁶ It may be protested that PA already contain symbols for conjunction and negation, say '&' and '¬'; so why do we adjoin new symbols ' \wedge ' and ' \sim '? The answer is that if P and Q are atomic formulae, then $P \& Q$ is *also* atomic in the present sense, since it is well-formed in PA; but $P \wedge Q$ is not. In order to be able to apply the previous theory, in which the conjunction of atomic formulae is not atomic, we need ' \wedge '. Nevertheless, for any $H \in K$ and atomic P and Q , $\varphi(P \& Q, H) = \varphi(P \wedge Q, H)$, so that confusion of '&' with ' \wedge ' causes no harm in practice. Similar remarks apply to negation, and to the provability interpretation of s4 in the next paragraph.

¹⁷ See J. C. C. McKinsey, 'On the Syntactical Construction of Systems of Modal Logic', *Journal of Symbolic Logic*, 10 (1945), 83-94.

PA, and whose well-formed formulae are formed out of the closed formulae of PA by use of the connectives $\&$, \neg , and \Box . (I say '&' and ' \neg ' to indicate that I am using the same conjunction and negation as in PA itself, not introducing new ones. See footnote 16, p. 71.) Then E is called a reflexive extension of PA iff: (1) It is an inessential extension of PA; (2) $\Box A$ is provable in E iff A is; (3) there is a valuation α , mapping the closed formulae of E into the set $\{T, F\}$, such that conjunction and negation obey the usual truth tables, all the true closed formulae of PA get the value T, $\alpha(\Box A) = T$ iff A is provable in E, and all the theorems of E get the value T. It can be shown that there are reflexive extensions of PA containing the axioms of S4 or even S4.1, but none containing S5.

Finally, using the usual mapping of intuitionistic logic into S4, we can get a model theory for the intuitionistic predicate calculus. We will not give this model theory here, but instead will mention, for propositional calculus only, a particular useful interpretation of intuitionistic logic that results from the model theory. Let E be any consistent extension of PA. We say a formula P of PA is *verified* in E iff it is provable in E. We take the closed wffs P of PA as atomic, and build formulae out of them using the intuitionistic connectives \wedge , \vee , \neg , and \supset . We then stipulate inductively: $A \wedge B$ is verified in E iff A and B are; $A \vee B$ is verified in E iff A or B is; $\neg A$ is verified in E iff there is no consistent extension of E verifying A ; $A \supset B$ is verified in E iff every consistent extension E' of E verifying A also verifies B .

Then every instance of a law of intuitionistic logic is verified in PA; but, e.g., $A \vee \neg A$ is not, if A is the Gödel undecidable formula. In future work, we will extend this interpretation further, and show that using it we can find an interpretation for Kreisel's system FC of absolutely free choice sequences.¹⁸ It is clear, incidentally, that PA can be replaced in the provability interpretations of S4 and S5 by any truth functional system (i.e., by any system whose models determine each closed formula as true or false); while the interpretation of intuitionism applies to any formal system whatsoever.

¹⁸ G. Kreisel, 'A Remark on Free Choice Sequences and the Topological Completeness' Proofs, *Journal of Symbolic Logic*, 23 (1958), 369-88.

VI

ESSENTIALISM AND QUANTIFIED MODAL LOGIC¹

TERENCE PARSONS

PROBLEMS involving essentialism are now receiving a great deal of attention from modal logicians and philosophers. Even a cursory glance at work in this field, however, soon reveals that there are many doctrines which go by this title. I will isolate and discuss one such doctrine. In particular, after isolating one version of essentialism (Sections I and II), I will argue that work in quantified modal logic can be and is independent of the acceptance of the truth of this doctrine (Sections III-V). In the last section (Section VI) I will attempt to show, on the basis of facts established in Sections III-V, just why this particular form of essentialism is a philosophically suspect doctrine. I will also argue that work in quantified modal logic need not even presuppose the *meaningfulness* of essentialist claims in any objectionable sense.

My arguments aim at (a) a clarification of one sort of essentialism, and (b) a partial vindication of quantified modal logic.

I. PRELIMINARY CLARIFICATION

To begin, let us dichotomize essentialist doctrines into two kinds. One kind has to do with what I shall call *individual* essences and the other with what I shall call *general* essences. The former doctrine makes some claim to the effect that some or all objects have characteristics (or properties) which are so intimately associated with the object that nothing else *could* (with emphasis on the 'could') have precisely those characteristics without being that object. This is meant to be a stronger thesis than the Identity of Indiscernibles, which holds merely that no two objects can simultaneously exist while sharing all properties. It is stronger in two ways: (1) it prohibits the simultaneous existence of two objects which share the same individual essence (even when they could differ in other of their properties), and (2) it makes a claim about what *might have been*: had

From *The Philosophical Review*, LXXVIII. 1 (January 1969), 35-52. Reprinted by permission of the author and *The Philosophical Review*.

¹ In addition to the authors cited in the paper, I am particularly indebted to John Vickers and to Kathryn Pyne Parsons for comments on earlier drafts, and to the referee of *The Philosophical Review* for help in improving the final draft.